

GIRSANOV THEOREM FOR FILTERED POISSON PROCESSES

L. DECREUSEFOND AND N. SAVY

ABSTRACT. Shot-noise and fractional Poisson processes are instances of filtered Poisson processes. We here prove Girsanov theorem for this kind of processes and give an application to an estimate problem.

1. INTRODUCTION

At the beginning of the theories of mathematical finance and performance analysis of telecommunications networks, asset prices or input traffics were modeled by Markovian diffusion processes. It is by now well known that these processes don't capture some of the essential features observed in the real datas. Namely, diffusion processes do not reflect long-range dependence exhibited in many traces [6, 7]. Alternative models are currently being considered such as Lévy processes, fractional Brownian motion, shot noise processes and different combinations of them. For instance, in [6], it is suggested that one could represent the evolution of an asset price by a process $S_t = \exp(B_t + N_t)$, where B is a standard Brownian motion and N is a shot noise process (for discussions about the validity of this model we refer to [6]). Motivated by this work, we here investigate some of the basic properties of processes which encompass shot-noise models. Given a marked Poisson process N and a deterministic kernel K , we define filtered Poisson process N^K , by

$$\begin{aligned} N_t^K &= \int_0^t K(t, s) dN_s \\ &= \int_0^t \int_E z K(t, s) \mu(ds, dz). \end{aligned}$$

If K is a convolution kernel, i.e., $K(t, s) = k(t - s)$ then N^K is usually called a shot-noise process. Shot-noise processes are used in numerous fields of applications, e.g. electronics, hydrology, climatology, telecommunications (see [9], [8], [11] and references therein), insurance (see [1] and references therein) and finance (see [6] and [10] for a review on this topic). It follows that filtered Poisson processes can, in the same fields, model more general phenomena so that we believe, it is an interesting class of processes to consider. This paper is organized as follows. In the next section, we give some

preliminaries on point processes. In section 3, we prove the Girsanov theorem for filtered Poisson processes and in section 4, we apply this result to an estimate problem.

2. PRELIMINARIES

Let $\mathbb{R}^d$ equipped with its borelian σ -algebra and $(\Omega, \mathfrak{F}, (\mathfrak{F}_t)_{t \in \mathbb{R}^+}, \mathbf{P})$ a filtered probability space. We consider that we are given an $\mathfrak{F}$ -adapted, marked Point process μ of intensity measure ν of the form (for details, see [3])

$$\nu(\omega, ds, dz) = \lambda(s) ds \eta(dz).$$

This means that there exist two sequences of random variables $(T_n)_{n \in \mathbb{N}^*}$ and $(Z_n)_{n \in \mathbb{N}^*}$ such that : $(T_n)_{n \in \mathbb{N}^*}$ is a strictly increasing sequence of $\mathbb{R}^+$ (jump times) verifying $\lim_{n \rightarrow \infty} T_n = +\infty$ and $(Z_n)_{n \in \mathbb{N}^*}$ is a sequence of $\mathbb{R}^d$ (the marks associated with the jump times), such that

$$\mu_{t,z} = \sum_{n \in \mathbb{N}^*} \delta_{(T_n, Z_n)}(t, z).$$

We associate with μ the Marked Poisson process N defined as:

$$N_t = \sum_{n \in \mathbb{N}^*} Z_n \mathbf{1}_{[T_n \leq t]}.$$

Hypothesis 1. *We assume now that our deterministic kernel satisfies a few regularity assumptions :*

- *K is triangular, i.e., $K(t, s) = 0$ for $s > t$.*
- *K is sufficiently integrable in the sense that $(s, z) \rightarrow zK(t, s)$ is in $\mathcal{L}^2(\nu)$ for any $t > 0$.*
- *K is continuously differentiable with respect to t and s in $\{(t, s) : 0 < s < t\}$.*

We now define a filtered Poisson process by

$$(1) \quad N_t^K = \int_0^t \int_E z K(t, s) \mu(ds, dz).$$

The first hypothesis thus appears as necessary if we want to have a non-anticipative process N^K . The other two are only technical assumptions. The next proposition is immediate.

Proposition 1. *The sample-paths of N^K are càdlàg iff $K(t, t)$ is finite for any t and that they are continuous iff $K(t, t) = 0$ for any t .*

3. GIRSANOV THEOREM

We hereafter assume that K is degenerate on the diagonal, that is to say $K(t, t) = 0$ for any t . It has thus a sense to look at the perturbations of the sample-paths which induce an absolutely continuous change of probability measure.

Theorem 1. *Let $\{\tilde{N}_t : t \geq 0\}$ the compensated Marked Poisson Process associated with N defined by*

$$\tilde{N}_t^K = \int_0^t \int_E z K(t, s) (\mu(ds, dz) - \nu(ds, dz)).$$

Consider a function $h \in \mathcal{L}^1(\nu)$ and introduce the probability measure $\mathbf{P}_h$ which is absolutely continuous with respect to $\mathbf{P}$ of Radon-Nikodym density given by:

$$\begin{aligned} Z_t &= \left. \frac{d\mathbf{P}_h}{d\mathbf{P}} \right|_{\mathfrak{F}_t} \\ (2) \quad &= \mathcal{ED} \left(\int_0^t \int_E h(s, z) (\mu - \nu)(ds, dz) \right)_t \end{aligned}$$

where $\mathcal{ED}(X)_t$ indicates the Doléans-Dade exponential of the process X at time t .

Let $N^{h,K}$ be defined by

$$N_t^{h,K} = \int_0^t \int_E z K(t, s) (\mu(ds, dz) - \nu(ds, dz)) - \int_0^t \int_E z K(t, s) h(s, z) \nu(ds, dz).$$

Then the following equality in law holds:

$$\mathcal{L}(\tilde{N}^K, \mathbf{P}_h) = \mathcal{L}(\tilde{N}^{h,K}, \mathbf{P})$$

Proof. Let $n \in \mathbb{N}^*$ and $(t_1, \dots, t_n) \in [0, T]^n$ fixed. When these instants are fixed, the processes

$$M_r^i : r \rightarrow \int_0^r \int_E z K(t_i, s) \mu(ds, dz)$$

are marked Point processes whose compensators are:

$$\Phi_r^i : r \rightarrow \int_0^r \int_E z K(t_i, s) \nu(ds, dz)$$

Hence, $(M^i - \Phi^i)_{i=1}^n$ is a $\mathbf{P}$ -martingale and the Girsanov theorem (see [3]) makes certain that:

$$\left(M^{i,h} : r \rightarrow M_r^i - \Phi_r^i - \int_0^r \frac{1}{Z_{s-}} d\langle M^i - \Phi^i, Z \rangle_s \right)_{i=1}^n$$

is a $\mathbf{P}_h$ -martingale. Z is defined by (2) so Z is solution of the equation:

$$R_t = 1 + \int_0^t \int_E R_{s-} [h(s, z) - 1] (\mu - \nu)(ds, dz)$$

Hence, we can write:

$$\begin{aligned}
(A) &= \langle M^i - \Phi^i, Z \rangle_s \\
&= \left\langle \int_0^\cdot \int_E zK(t_i, s)(\mu - \nu)(ds, dz), 1 + \int_0^\cdot \int_E Z_{s-}[h(s, z) - 1](\mu - \nu)(ds, dz) \right\rangle_s \\
&= \left\langle \int_0^\cdot \int_E zK(t_i, s)(\mu - \nu)(ds, dz), \int_0^\cdot \int_E Z_{s-}[h(s, z) - 1](\mu - \nu)(ds, dz) \right\rangle_s \\
&= \int_0^s zK(t_i, s)Z_{s-}[h(s, z) - 1]d \left\langle \int_0^\cdot \int_E (\mu - \nu)(ds, dz); \int_0^\cdot \int_E (\mu - \nu)(ds, dz) \right\rangle_s \\
&= \int_0^s \int_E zK(t_i, s)Z_{s-}[h(s, z) - 1]\nu(ds, dz)
\end{aligned}$$

We can conclude that:

$$\begin{aligned}
M_r^{i,h} &= M_r^i - \Phi_r^i - \int_0^r \frac{1}{Z_{s-}} d \langle M^i - \Phi^i, Z \rangle_s \\
&= M_r^i - \Phi_r^i - \int_0^r \frac{1}{Z_{s-}} \int_E zK(t_i, s)Z_{s-}[h(s, z) - 1]\nu(ds, dz) dz \\
&= M_r^i - \Phi_r^i - \int_0^r \int_E zK(t_i, s)[h(s, z) - 1]\nu(ds, dz) \\
&= \int_0^r \int_E zK(t_i, s)\mu(ds, dz) - \Phi_r^i - \int_0^r \int_E zK(t_i, s)h(s, z)\nu(ds, dz) + \Phi_r^i \\
(3) \quad &= \int_0^r \int_E zK(t_i, s)\mu(ds, dz) - \int_0^r \int_E zK(t_i, s)h(s, z)\nu(ds, dz)
\end{aligned}$$

thus,

$$r \rightarrow \int_0^r \int_E zK(t_i, s)\mu(ds, dz) - \int_0^r \int_E zK(t_i, s)h(s, z)\nu(ds, dz)$$

is a $\mathbf{P}_h$ -martingale and

$$(4) \quad r \rightarrow \int_0^r \int_E zK(t_i, s)h(s, z)\nu(ds, dz)$$

is the $\mathbf{P}_h$ -compensator of M^i .

>From (3), $M^{i,h}$ is written:

$$M^{i,h} : r \rightarrow \int_0^r \int_E zK(t_i, s)(\mu(ds, dz) - h(s, z)\nu(ds, dz))$$

So its compensator under $\mathbf{P}$ is (4).

It follows (see [3]) that:

$$\mathcal{L} \left([r \rightarrow M_r^i]_{i=1}^n ; \mathbf{P}_h \right) = \mathcal{L} \left([r \rightarrow M_r^{i,h}]_{i=1}^n ; \mathbf{P} \right)$$

Now, taking $r = \sup_{1 \leq i \leq n} t_i$, the kernel being triangular, the result holds. $\square$

4. AN APPLICATION

Suppose now that we are given to observe the paths of a perturbed filtered Poisson process:

$$X_t^\theta = \tilde{N}_t^K - \theta t \text{ with } \theta > 0.$$

We now investigate the problem of the estimate of θ . Such a problem has been thoroughly investigated when a standard Brownian motion is put in place of N (i.e., a fractional Brownian motion is substituted to $\tilde{N}^K$) [2, 4, 5]. We here need to assume that ν and K are such that there exists one and only one function ϕ such that:

$$\int_0^t \int_E z K(t, s) \phi(s) \nu(ds, dz) = t.$$

We then introduce the likelihood function:

$$Z_t^\theta = \left. \frac{d\mathbf{P}_\theta}{d\mathbf{P}} \right|_{\mathfrak{F}_t} = \mathcal{ED} \left(\int_0^\cdot \int_E \theta \phi(s) (\mu - \nu)(ds, dz) \right)_t.$$

According to the previous theorem, the processes X^θ under $\mathbf{P}$ and $\tilde{N}^K$ under $\mathbf{P}_\theta$ have the same law thus an estimate of θ is given by:

$$\hat{\theta}_t = \underset{\theta \in [0,1]}{\operatorname{Argmax}} Z_t^\theta.$$

Unfortunately, the exact expression of the Doléans-Dade exponential is so intricate in case of jump processes that we have to find another expression of Z^θ more suitable for computations $\hat{\theta}_t$.

Lemma 1. *If $\phi \in \mathcal{L}^1(\nu)$ and $\ln(1 + \theta\phi) \in \mathcal{L}^1(\nu)$ then:*

$$\left\{ \mathcal{ED} \left(\int_0^\cdot \int_E \theta \phi(s) (\mu - \nu)(ds, dz) \right)_t : 0 \leq t \leq T \right\} = \left\{ \exp(Y_t^\theta) : 0 \leq t \leq T \right\}$$

with for any $0 \leq t \leq T$:

$$Y_t^\theta = \int_0^t \int_E \ln(1 + \theta\phi(s)) \mu(ds, dz) - \int_0^t \int_E \theta\phi(s) \nu(ds, dz)$$

Proof. Apply Itô formula to the function $x \rightarrow \exp(x)$ with respect to the jump process Y^θ . One gets:

$$\begin{aligned}
& \exp(Y_t^\theta) - 1 \\
&= \int_0^t \exp(Y_{s-}^\theta) dY_s^\theta + \sum_{s \leq t} [\exp(Y_s^\theta) - \exp(Y_{s-}^\theta) - \exp(Y_{s-}^\theta)(Y_s^\theta - Y_{s-}^\theta)] \\
&= \int_0^t \exp(Y_{s-}^\theta) dY_s^\theta + \int_0^t \int_E [\exp(Y_{s-}^\theta + \ln(1 + \theta\phi(s))) - \exp(Y_{s-}^\theta) \\
&\quad - \exp(Y_{s-}^\theta)(\ln(1 + \theta\phi(s)))] \mu(ds, dz) \\
&= \int_0^t \exp(Y_{s-}^\theta) dY_s^\theta + \int_0^t \int_E [\exp(Y_{s-}^\theta + \ln(1 + \theta\phi(s))) \\
&\quad - \exp(Y_{s-}^\theta)(1 + \ln(1 + \theta\phi(s)))] \mu(ds, dz)
\end{aligned}$$

But we have:

$$dY_s^\theta = \int_E \ln(1 + \theta\phi(s)) \mu(ds, dz) - \int_E \theta\phi(s) \nu(ds, dz)$$

hence:

$$\begin{aligned}
& \exp(Y_t^\theta) - 1 \\
&= \int_0^t \int_E \exp(Y_{s-}^\theta) [\ln(1 + \theta\phi(s)) \mu(ds, dz) - \theta\phi(s) \nu(ds, dz)] \\
&\quad + \int_0^t \int_E [\exp(Y_{s-}^\theta + \ln(1 + \theta\phi(s))) \\
&\quad - \exp(Y_{s-}^\theta)(1 + \ln(1 + \theta\phi(s)))] \mu(ds, dz) \\
&= \int_0^t \int_E \exp(Y_{s-}^\theta) [\ln(1 + \theta\phi(s)) - (1 + \ln(1 + \theta\phi(s)))] \\
&\quad + \exp(\ln(1 + \theta\phi(s)))] \mu(ds, dz) \\
&\quad - \int_0^t \int_E \exp(Y_{s-}^\theta) \theta\phi(s) \nu(ds, dz) \\
&= \int_0^t \int_E \exp(Y_{s-}^\theta) \theta\phi(s) \mu(ds, dz) - \int_0^t \int_E \exp(Y_{s-}^\theta) \theta\phi(s) \nu(ds, dz) \\
&= \int_0^t \int_E \exp(Y_{s-}^\theta) \theta\phi(s) (\mu - \nu)(ds, dz)
\end{aligned}$$

It follows that the process $\exp(Y^\theta)$ is a solution to the Stochastic Differential Equation which defines the Doléans-Dade exponential. By uniqueness of the solution, the result holds. $\square$

Proposition 1. Suppose $\phi \in \mathcal{L}^1(\nu)$ and $\ln(1 + \theta\phi) \in \mathcal{L}^1(\nu)$. Then, for any $t \in [0, T]$, $\hat{\theta}_t$ exists and is the unique positive solution of

$$(5) \quad \sum_{T_j \leq t} \frac{\phi(T_j)}{1 + \hat{\theta}_t \phi(T_j)} = \int_0^t \phi(s) \lambda(s) ds.$$

Proof. Fix $t \in [0, T]$. The likelihood function reaches its maximum when the function:

$$f : \theta \rightarrow \int_0^t \int_E \ln(1 + \theta\phi(s)) \mu(ds, dz) - \int_0^t \int_E \theta\phi(s) \nu(ds, dz)$$

reaches its maximum. Noticing that:

$$\begin{aligned} f'(\theta) &= \int_0^t \int_E \frac{\phi(s)}{1 + \theta\phi(s)} \mu(ds, dz) - \int_0^t \int_E \phi(s) \nu(ds, dz) \\ f''(\theta) &= \int_0^t \int_E \frac{-\phi(s)^2}{[1 + \theta\phi(s)]^2} \mu(ds, dz), \end{aligned}$$

we see that the function f is concave hence that it admits a unique maximum and that this maximum $\hat{\theta}_t$ satisfies (5). $\square$

Remark 1. If we consider the fractional Poisson Process: this means that K is taken to be the kernel K_H associated to fractional Brownian motion (see [2]):

$$K_H(t, s) = \frac{1}{\Gamma(H + 1/2)} (t - s)^{H-1/2} F(H - 1/2, 1/2 - H, H + 1/2, 1 - t/s),$$

where $F(a, b, c, z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 u^{b-1} (1-u)^{c-b-1} (1-zu)^{-a} du$, and that N is a Poisson process with constant intensity λ . Then, we know (see Decreusefond Üstünel [2]) that ϕ is given by:

$$(6) \quad s \rightarrow \phi(s) = \frac{\Gamma(\frac{3}{2} - H)}{\Gamma(2 - 2H)} \frac{s^{\frac{1}{2}-H}}{\lambda}$$

The theorem can be applied because $\phi \in \mathcal{L}^1(\nu)$ as soon as $\frac{1}{2} < H < 1$ and in this case so does $\ln(1 + \theta\phi)$.

We now prove that for $\theta > 0$, $\hat{\theta}_t$ is strongly consistant, i.e., that $\hat{\theta}_t$ converges a.s. to θ .

Lemma 2. The process $\hat{\theta}_t$ is decreasing on any interval $[T_n, T_{n+1}[$.

Proof. Let $\psi(x, y) = x/(1 + xy)$, it is clear that for $x > 0$, the partial map $(y \mapsto \psi(x, y))$ is decreasing. We know that $\hat{\theta}_t$ is the solution of the equation

$$(7) \quad \sum_{T_j \leq t} \psi(\phi(T_j), \hat{\theta}_t) = \int_0^t \phi(s) \lambda(s) ds.$$

For $t \in [T_n, T_{n+1}[$, the number of terms in the left-hand-side in (7) is constant and the right-hand-side is an increasing function of t . It follows that $\hat{\theta}_t$ must decrease between T_n and T_{n+1} . $\square$

To go further, we need additional hypothesis on ϕ . Note that all these hypothesis are satisfied by ϕ as defined in (6). We don't know whether $\hat{\theta}_{T_n}$ is decreasing sequence and we cannot thus conclude to the convergence of $\hat{\theta}_t$. However, we now prove that it is a bounded process.

Lemma 3. *Assume $\theta > 0$ and that ϕ and $\ln(1+\theta\phi)$ belong to $L^1([0, T], \lambda(s)ds)$ for any $T > 0$. Assume that*

$$\lim_{t \rightarrow \infty} \int_0^t \phi^2(s) \lambda(s) ds = \infty$$

and that

$$(8) \quad \lim_{t \rightarrow \infty} \frac{\int_0^t \phi^{2+j}(s) \lambda(s) ds}{\int_0^t \phi^2(s) \lambda(s) ds} = 0, \text{ for any } j > 0.$$

Then, $\{\hat{\theta}_t, t \geq 0\}$ is $\mathbf{P}_\theta$ -a.s. bounded.

Proof. Let $M > \theta$ and consider

$$A_M = \{\omega \in \Omega, \limsup_{n \rightarrow \infty} \hat{\theta}_t(\omega) \geq M\}.$$

On A_M , there exists a sequence $\{t_n, n \geq 1\}$ of positive reals such that $\hat{\theta}_{t_n} \geq M$ for any $n \geq 1$. Thus

$$\sum_{T_j \leq t_n} \psi(\phi(T_j), \hat{\theta}_{t_n}) \leq \sum_{T_j \leq t_n} \psi(\phi(T_j), M)$$

By the very definition of $\hat{\theta}_t$,

$$\sum_{T_j \leq t_n} \psi(\phi(T_j), \hat{\theta}_{t_n}) = \int_0^{t_n} \phi(s) \lambda(s) ds,$$

thus

$$(9) \quad \begin{aligned} & \int_0^{t_n} \phi(s) \lambda(s) ds - \int_0^{t_n} \psi(\phi(s), M)(1 + \theta\phi(s)) \lambda(s) ds \\ & \leq \sum_{T_j \leq t_n} \psi(\phi(T_j), M) - \int_0^{t_n} \psi(\phi(s), M)(1 + \theta\phi(s)) \lambda(s) ds. \end{aligned}$$

Left-hand-side of Equation (9) can be simplified as

$$\begin{aligned} & \int_0^{t_n} \phi(s) \lambda(s) ds - \int_0^{t_n} \psi(\phi(s), M)(1 + \theta\phi(s)) \lambda(s) ds \\ & = (M - \theta) \int_0^{t_n} \frac{\phi^2(s) \lambda(s)}{1 + M\phi(s)} ds. \end{aligned}$$

Thus, Equation (9) now reads as

$$(10) \quad (M - \theta) \int_0^{t_n} \frac{\phi^2(s)\lambda(s)}{1 + M\phi(s)} ds \leq \sum_{T_j \leq t_n} \psi(\phi(T_j), M) - \int_0^{t_n} \psi(\phi(s), M)(1 + \theta\phi(s))\lambda(s) ds.$$

Furthermore, since $\phi \geq 0$ and $M > \theta$, we have

$$(11) \quad \int_0^t \frac{\phi^2(s)\lambda(s)(1 + \theta\phi(s))}{(1 + M\phi(s))^2} ds \leq \int_0^t \frac{\phi^2(s)\lambda(s)}{1 + M\phi(s)} ds.$$

On the other hand, Central Limit theorem for martingale, says that, $\mathbf{P}_\theta$ a.s.,

$$\frac{\sum_{T_j \leq t_n} \psi(\phi(T_j), M) - \int_0^{t_n} \psi(\phi(s), M)(1 + \theta\phi(s))\lambda(s) ds}{\int_0^{t_n} \psi(\phi(s), M)^2 \lambda(s)(1 + \theta\phi(s)) ds} \xrightarrow{n \rightarrow \infty} 0,$$

Then, divide both sides of (10) by $\int_0^{t_n} \psi(\phi(s), M)\phi(s)\lambda(s) ds$ and let n go to infinity. In virtue of (8) and (11), this yields to the contradiction that on A_M , $\theta \geq M$. Finally, $\mathbf{P}_\theta(A_M) = 0$ and thus $\{\hat{\theta}_t, t > 0\}$ is $\mathbf{P}_\theta$ -a.s. bounded. $\square$

Theorem 2. Assume $\theta > 0$ and that ϕ and $\ln(1 + \theta\phi)$ belong to $L^1([0, T], \lambda(s)ds)$ for any $T > 0$. Assume that

$$\lim_{t \rightarrow \infty} \int_0^t \phi^2(s)\lambda(s) ds = \infty$$

and that

$$\lim_{t \rightarrow \infty} \frac{\int_0^t \phi^{2+j}(s)\lambda(s) ds}{\int_0^t \phi^2(s)\lambda(s) ds} = 0, \text{ for any } j > 0.$$

Then $\hat{\theta}_t$ tends P_θ -a.s. to θ .

Proof. Let $N_t(\phi) = \sum_{T_n \leq t} \phi(T_n)$. According to (7), we have

$$\begin{aligned} 0 &= \sum_{T_n \leq t} \frac{\phi(T_n)}{1 + \hat{\theta}_t \phi(T_n)} - \int_0^t \phi(s)\lambda(s) ds \\ &= N_t(\phi) - \int_0^t \phi(s)\lambda(s)(1 + \theta\phi(s)) ds \\ &\quad - \hat{\theta}_t \left(N_t(\phi^2) - \int_0^t \phi^2(s)\lambda(s)(1 + \theta\phi(s)) ds \right) \\ &\quad + \hat{\theta}_t^2 \sum_{T_n \leq t} \frac{\phi(T_n)^3}{1 + \hat{\theta}_t \phi(T_n)} - \theta \hat{\theta}_t \int_0^t \phi^3(s)\lambda(s) ds \\ &\quad + (\theta - \hat{\theta}_t) \int_0^t \phi^2(s)\lambda(s) ds. \end{aligned}$$

After some simple algebra, this yields to

$$\begin{aligned}
(12) \quad & (\theta - \hat{\theta}_t) \left(\int_0^t \phi^2(s) \lambda(s) ds + \hat{\theta}_t \int_0^t \phi^3(s) \lambda(s) ds \right) \\
&= - \left(N_t(\phi) - \int_0^t \phi(s) \lambda(s) (1 + \theta \phi(s)) ds \right) \\
&+ \hat{\theta}_t \left(N_t(\phi^2) - \int_0^t \phi^2(s) \lambda(s) (1 + \theta \phi(s)) ds \right) \\
&- \hat{\theta}_t^2 \left(N_t(\phi^3) - \int_0^t \phi^3(s) \lambda(s) (1 + \theta \phi(s)) ds \right) \\
&- \hat{\theta}_t^2 \sum_{T_n \leq t} \left(\frac{\phi(T_n)^3}{1 + \hat{\theta}_t \phi(T_n)} - \phi(T_n)^3 \right) + \theta \hat{\theta}_t^2 \int_0^t \phi^4(s) \lambda(s) ds.
\end{aligned}$$

For any deterministic ζ , the process $\{N_t(\zeta) - \int_0^t \zeta(s) \lambda(s) (1 + \theta \phi(s)) ds, t \geq 0\}$ is a martingale whose square bracket given by

$$\int_0^t \zeta(s)^2 \lambda(s) (1 + \theta \phi(s)) ds.$$

It follows from hypothesis and Central Limit Theorem for martingales that

$$(13) \quad \lim_{t \rightarrow \infty} \frac{N_t(\phi^j) - \int_0^t \phi(s)^j \lambda(s) (1 + \theta \phi(s)) ds}{\int_0^t \phi^2(s) \lambda(s) ds} = 0,$$

for any $j \geq 1$. Moreover,

$$(14) \quad \sum_{T_n \leq t} \left(\frac{\phi(T_n)^3}{1 + \hat{\theta}_t \phi(T_n)} - \phi(T_n)^3 \right) = \hat{\theta}_t \sum_{T_n \leq t} \frac{\phi(T_n)^4}{1 + \hat{\theta}_t \phi(T_n)}.$$

Thus,

$$0 \leq \sum_{T_n \leq t} \left(\frac{\phi(T_n)^3}{1 + \hat{\theta}_t \phi(T_n)} - \phi(T_n)^3 \right) \leq \hat{\theta}_t N_t(\phi^4).$$

According to (13) and to hypothesis (8), we have

$$(15) \quad \lim_{t \rightarrow \infty} \frac{N_t(\phi^4)}{\int_0^t \phi^2(s) \lambda(s) ds} = 0, \quad \mathbf{P}_\theta \text{ a.s.}$$

Divide (12) by $\int_0^t \phi^2(s) \lambda(s) ds$ and let t goes to infinity, it follows from (13) and (15) that $\hat{\theta}_t$ converges to θ . $\square$

REFERENCES

- [1] A. Dassios and J.W. Jang. The shot-noise process and its distribution at claim jump times as the intensity of a doubly stochastic proces. Preprint.
- [2] L. Decreusefond and A.S. Üstünel. Stochastic analysis of the fractional Brownian motion. *Potential Anal.*, 10(2):177–214, 1999.
- [3] J. Jacod. *Calcul stochastique et problèmes de martingales*, volume 714 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979.

- [4] M. L. Kleptsyna and A. Le Breton. A Cameron-Martin type formula for general Gaussian processes—a filtering approach. *Stoch. Stoch. Rep.*, 72(3-4):229–250, 2002.
- [5] M. L. Kleptsyna, A. Le Breton, and M.-C. Roubaud. General approach to filtering with fractional Brownian noises—application to linear systems. *Stochastics Stochastics Rep.*, 71(1-2):119–140, 2000.
- [6] C. Kühn. *Shocks and Choices - an Analysis of Incomplete Market Models*. PhD thesis, Technische Universität München, 2002.
- [7] W.E Leland, M.S. Taqqu, W. Willinger, and D.V. Wilson. On the self-similar nature of ethernet traffic. *IEEE/ACM trans. Networking*, 2(1):1–15, Februar 1994.
- [8] D.L Snyder M.I Miller. *Random Point Processes in Time and Space*. Springer Verlag, New York, 1991.
- [9] E. Parzen. *Stochastic processes*, volume 24 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1999. Reprint of the 1962 original.
- [10] G. Samorodnitsky. A class of shot noise models for financial applications. In *Athens Conference on Applied Probability and Time Series Analysis, Vol. I (1995)*, volume 114 of *Lecture Notes in Statist.*, pages 332–353. Springer, New York, 1996.
- [11] S. Yue and M. Hishino. The general cumulants for a filtered point processes. *Applied Mathematical Modelling*, 25:193–201, 2001.

DÉPARTEMENT INFORMATIQUE ET RÉSEAUX, ÉCOLE NATIONALE SUPÉRIEURE DES
TÉLÉCOMMUNICATIONS, 46, RUE BARRAULT, 75634 PARIS CEDEX 13, FRANCE

INSTITUT DE RECHERCHE MATHÉMATIQUES DE RENNES, UNIVERSITÉ DE RENNES 1,
35042 RENNES CEDEX, FRANCE